



Neonatal On-line Education Learning Objectives



By the end of the webinar, the learner will be able to:

Webinar 1: Living a Great Story

- Objectives:
1. Name two possible neurodevelopmental outcomes of prematurity.
 2. Identify the purpose of the Just Be step in caregiving.

Webinar 2: Brain & Sensory Development: Clinical Application in the NICU

- Objectives:
1. Recognize two neurosensory vulnerabilities of the preterm and critically ill, hospitalized infant.
 2. List three methods to adapt care that positively support the ongoing development of the brain and sensory system.

Webinar 3: Pain: An Undeniable Indicator for Caregiver Intervention

- Objectives:
1. Identify two potential long-term effects of unmanaged pain in neonates.
 2. Name three pain management interventions/techniques to use in the NICU.

Webinar 4: Positioning & Handling of the Neonate

- Objectives:
1. Recognize a common postural issue of prematurity and how positioning prevents this issue.
 2. List three functional activities affected by positioning.

Webinar 5: Infant-Driven Caregiving

- Objectives:
1. Identify four goals of Infant-Driven Caregiving.
 2. List three activities where 2-person caregiving is beneficial.

Webinar 6: Inside the NICU Parent Experience: Meeting Parents Where They Are

- Objectives:
1. Recognize the traumatic experience of having a critically ill child in the NICU
 2. Recognize the traumatic experience of having a critically ill child in the NICU.
 3. Identify two strategies to meet the psychosocial needs of NICU parents

Webinar 7: Nurturing Care of Mother-Infant Dyads Impacted by Opioid Use

- Objectives:
1. Identify two global and national trends in opioid use.
 2. Identify three non-pharmacologic interventions for management of Neonatal Abstinence Syndrome.

Webinar 8: Environment of Care

- Objectives:
1. List three environmental modifications that can be made in the NICU to improve infant neurodevelopmental outcomes.
 2. Recognize three research articles that highlight non-pharmacological approaches to reduce pain in the NICU.

